


Quarantine/Isolation Site Discharge Planning

Patient Name: _____

DOB: _____

Room: _____

Admission Date: _____



Medically Cleared for Discharge by:

Provider Name _____

Provider Signature _____

Date & Time _____

1. For patients returning to congregate setting, test pt on day 4 of the exposure if no previous test(s) were (-). Quarantine patients who tested negative need to quarantine for 10 days regardless of symptoms.
2. For patients in continued close contact with a COVID (+) person, the last close contact date is the date the (+) person completes their isolation.
3. Isolate 20 days for patients who have severe disease or severely immunocompromised.
 Severe disease: Individuals who have respiratory failure requiring nasal cannula greater than 6L/minute, high flow nasal cannula/BiPAP/mechanical ventilation or ECMO, or lung infiltrates >50% on chest X-ray or CT scan, or other manifestations such as clotting, etc., septic shock, and/or multiple organ dysfunction.
 Severely Immunocompromised: chemotherapy for cancer, untreated HIV infection with CD4 T lymphocyte count <200/mm³, combined primary immunodeficiency disorder, and receipt of prednisone >20mg/day for more than 14 days.
 Diabetes, end stage renal disease, and advanced age are not considered severely immunocompromising
4. If patient becomes symptomatic, restart their isolation period with the symptom onset date.
5. The medical clearance date is the date AFTER patient completes their quarantine/isolation. For example, if the symptom onset date is 7/12, then the clearance date is 7/23.